
Treatment Practices In Patients With VTE - Survey Results


QUESTIONS #1






Treatment Practices In Patients With VTE - Survey Results


QUESTION #3






Treatment Practices In Patients With VTE - Survey Results

QUESTION #5






Treatment Practices In Patients With VTE – Survey Results

QUESTION #7


7. Inpatient: Which activity level do you order?		Create Chart	Create Chart 🕴 Download	
		Response Percent	Response Count	
Bed rest		4.7%	8	
Chair	I	0.6%	1	
Bathroom privileges only	_	14.8%	25	
Free Ambulation		79.9%	135	
		answered question	169	
		skipped question	21	



Treatment Practices In Patients With VTE – Survey Results


QUESTION #9

9. What do you routinely recommend?	Create Chart	♦ Download
	Response Percent	Response Count
Thrombo-embolic deterrent stockings (TED hose)	9.5%	14
20/30 stockings	48.0%	71
30/40 stockings	39.9%	59
Ace wraps	2.7%	4
	answered question	148
	skipped question	42





Treatment Practices In Patients With VTE - Survey Results

QUESTION #11



QUESTION #12





Treatment Practices In Patients With VTE - Survey Results

QUESTION #14

14. What do you routinely recommend?	Create Chart	♦ Download
	Response Percent	Response Count
Thrombo-embolic deterrent stockings (TED hose)	6.3%	9
20/30 stockings	55.9%	80
30/40 stockings	36.4%	52
Ace wraps	1.4%	2
	answered question	143
	skipped question	47

